PM10 & Childhood Asthma in Maricopa County

- EPA challenge grant 2006
- The impact of air pollution on children's health: the latest medical research


Asthma in Arizona

- Asthma prevalence above national average
- Up from 11% in 2000 to 15% in 2009
- 700,000+ Arizonans have been told by their doctor they have asthma
- Ambulatory-sensitive disease:
 with good primary care &
 avoidance of triggers, ER &
 hospital visits can be prevented





Asthma in Arizona



- AHCCCS paid 50% of all asthma ER visits, Medicare 6.7% = 56.7% public \$\$
- Total cost:\$44.5 million
- AHCCCS paid 39.1%
 & Medicare paid
 23.8% of the hospital
 bills = 62.9% public
 \$\$

Goals of the Study

- Exchange quality-assured hospital & ER admissions data & air quality data with partners
- Examine the correlations between data sets




Photo courtesy of azdeq.gov



Goals of the Study, cont.

 Develop a predictive model for asthma episodes & air quality thresholds in Maricopa County

 Investigate possibility of creating an alert system on the potential health effects of poor air

Partners







- ADEQ provided air monitor data
- ADHS provided data on ER visits & hospitalizations for asthma as a primary diagnosis
- ASU analyzed the data & provided predictive modeling for weather & pollution



Study Design

- Area: 168 census tracts in Metro Phoenix
- Boundary: Dunlap Rd., 52nd St., Eliot Rd & 75th Ave.
- Each census tract in area was located within 5 miles of a permanent PM₁₀ monitor
- Mapped PM₁₀ concentrations via monitoring & modeling



Central Phoenix Continuous PM10 Monitoring Sites

Monitors

Permanent:

CP - Central Phoenix

DC - Durango Complex

WF - West 43rd Avenue

WP - West Phoenix

GR - Greenwood

SP - South Phoenix

SS1 - Supersite

Temporary:

MRV - Maryvale

VGC - Valley Garden

Center

WVR - Weaver's Auto

Service

CSA - Community

Service






Photo courtesy of azdeq.gov

Study Design

- Analyzed 5,000+ hospitalizations & ER visits for asthma as the 1st diagnosis
- Population: children age 5-18
- Duration: 21 months, from January 1, 2005 to September 30, 2006
- Controls: reviewed 24-hour avg. PM₁₀ from 1-7 days before the health event in each case
- Modeled weather & pollution predictions for potential new warning systems

Results

Positive correlations between elevated PM₁₀ & high asthma incidence in Metro Phoenix




Photo courtesy of niehs.nih.gov

Results, cont.

- 13.7% increase in asthma events when PM_{10} rose by 36.4 $\mu g/m3$ (in range of 30-56 $\mu g/m3$)
- This is a change in the daily average PM_{10} from the 25th percentile to the 75th percentile
- Statistically significant at 95% confidence level

Asthma ER & hospital visits linked to elevated PM10





Conclusions



Photo courtesy of gnews.com

- This is a stronger effect than seen in other studies where such site-specific health and air quality data may not have been available
 - Significant increase in the likelihood that children may have an asthma event with a relatively small increase in PM₁₀

Study Conclusions

- The at-risk population is greater than previously observed
- Sensitive populations need greater protections than the current systems in place
- The study design can be used to model other pollutants such as PM_{2.5}, ozone, or a combination of pollutants



Photo courtesy of coyoteblog.com

Research on air pollution and asthma

- PM_{2.5}: Utah Valley Steel Mill closed for 13 mo. & reopened (1980s)
 - 89% increase in childhood hospital admissions during months mill was open
- Ozone: 1996 Atlanta Olympic Games, city traffic curtailed for 3 weeks
 - Peak ozone fell by 28%, Medicaid claims for asthma dropped 48%

Traffic-generated air pollution

- CA: longitudinal study in 12 communities-urban, suburban & rural
 - Showed reduced lung function growth linked to PM₁₀ concentration near urban highways
 - Long term exposure associated with changes in lung function in teens and adults
 - Forced respiratory volume (FEV) was lower in people who live in polluted areas

Traffic-generated pollution

- NJ & PA study of pregnant mothers who lived near highway toll plazas
 - After E-Z pass system introduced, traffic passed through tolls faster, no idling
 - Lower pollution associated with 10.8% reduced prematurity and
 - Low birthweight reduced by 11.8%

Effects of Inhaled PM:

• **Lungs**: *inflammation*, oxidative stress, alters pulmonary reflexes, reduces lung function

 Blood vessels: inflammation of lining, accelerates atherosclerosis, vasoconstriction & hypertension



REE Effects of Inhaled PM:

 Heart: abnormal rhythm, increases ischemia, exacerbates heart failure

 Blood: activates white blood cells, alters flow, increases coagulation, reduces oxygen saturation

• Brain: reduces blood flow (ischemia)



- Asthma incidence may be related to air pollution exposure in utero and early life
- 1st trimester exposure to PM₁₀ and 2nd trimester NO₂ exposures associated with lower lung function in children age 6-11
- Ozone, PM_{2,5} & CO are harmful to infants at levels below the EPA thresholds



Photo courtesy of sbcapcd.org

REUSABLE TOTE BAGS
HERVE ELECTRICITY MORE
HER REFUEL AFTER DARK
BIKE MORE RAKE MORE

improvements in air quality have bigger improvements in health, measured by life expectancy?

Studied 51 metro areas from 1980-2000

 YES: a 10 mcg/m³ reduction in PM_{2.5} is associated with 1 year increase in life expectancy



Policy implications

- Air pollution is recognized as a serious public health hazard
- Adverse effects on respiratory system occur at levels below the current ambient standards
- Focus control strategies on specific sources & constituents that are damaging to health

Policy implications



Photo courtesy of ehow.com



Photo courtesy of examiner.com

- Change zoning for schools & homes to increase distance from busy roadways
- Develop alert systems so atrisk people can decrease exposure on bad air days
- Study children's environmental health longterm
- Current regulations could be more stringent

Sources

- Protecting our Children: Assessing the Link Between PM₁₀
 Pollution and Childhood Asthma In Maricopa County.
 Children's Health Challenge Grant Project, December 2008.
 ADEQ.
 www.adea.gov/ceh/download/Exchange%20Grant%20Sum
 - <u>www.adeq.gov/ceh/download/Exchange%20Grant%20Sum</u> <u>mary.pdf</u>
- J. Currie, SH Ray and M Needle: Quasi-Experimental Studies Suggest that Lowering Air Pollution Benefits Infants and Children's Health. Health Affairs 30, #12:2391-2399. 2011.
- J Grigg: Air pollution and children's respiratory health gaps in the global evidence. Clinical and Experimental Allergy 41:1072-1075. 2011.

Sources

- DA Searing & N Rabinovitch. Environmental pollution and lung effects in children. Current Opinion in Pediatrics 23:314-318. 2011
- CA Pope, M Ezzati & D Dockery. Fine-Particulate Air Pollution and Life Expectancy in the United States. New England Journal of Med. Jan. 22, 2009
- Arizona Asthma Coalition:
 Barbara Burkholder barbburk5@msn.com